

Ab Initio Calculations of the Chiroptical Properties of Z-Configured Polymethine Dyes

V. Buß, M. Schreiber

published in

Modern Methods and Algorithms of Quantum Chemistry,
J. Grotendorst (Ed.), John von Neumann Institute for Computing,
Jülich, NIC Series, Vol. 2, ISBN 3-00-005746-3, p. 33, 2000.

© 2000 by John von Neumann Institute for Computing

Permission to make digital or hard copies of portions of this work for personal or classroom use is granted provided that the copies are not made or distributed for profit or commercial advantage and that copies bear this notice and the full citation on the first page. To copy otherwise requires prior specific permission by the publisher mentioned above.

<http://www.fz-juelich.de/nic-series/>

Ab initio Calculations of the Chiroptical Properties of Z-Configured Polymethine Dyes


V. Buß, M. Schreiber

Institut für Physikalische und Theoretische Chemie

Gerhard-Mercator-Universität Duisburg

Postfach, D-47048 Duisburg

Ab initio calculations of a series of inherently chiral, all-Z-configured polymethine dyes



are presented. The experimentally known chiroptical properties of the monomethine **1** and trimethine **2** make these compounds and their homologue **3** ideally suited to test the applicability of quantum mechanical calculations.

In compound **1** the chromophore is forced into a twisted all-Z-conformation by steric interaction of the end groups, while in **2** the presence of a t-butyl group in the *meso*-position enforces this conformation, which is manifest in the UV/Vis spectra not only in the reduced intensity of the longest wavelength absorption (methine band), but also in the occurrence of a „cis-peak“ at shorter wavelengths^[1].

Using the *MOLCAS* program-package excited states were calculated with the *CASSCF* and *CASPT2* methods. Electric and magnetic dipole transition moments were calculated, yielding UV/Vis and CD-spectra in good agreement with experimental results. The negative sign of the CD-band observed for P-helical monomethine **1** which appears to violate established helicity rules can be understood with the aid of a component analysis^[2,3] and was confirmed by the quantitative calculations. Additional calculations on the as yet not synthesized pentamethine **3** allow to predict the theoretical spectra of this compound.

[1] L. Zechmeister, Chem. Rev. 34 (1944) 267

[2] L. Eggers, V. Buß, G. Henkel, Angew. Chem. 108 (1996) 973

[3] L. Eggers, K. Kolster, V. Buß, Chirality 9 (1997) 243